

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Robert Katz Publications

Research Papers in Physics and Astronomy

March 1952

Radioactivity of Eu^{152,154}

Robert Katz

University of Nebraska-Lincoln, rkatz2@unl.edu

Milford R. Lee

Kansas State College, Manhattan, Kansas

Follow this and additional works at: <https://digitalcommons.unl.edu/physicskatz>



Part of the [Physics Commons](#)

Katz, Robert and Lee, Milford R., "Radioactivity of Eu^{152,154}" (1952). *Robert Katz Publications*. 21.
<https://digitalcommons.unl.edu/physicskatz/21>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Robert Katz Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Physical Review 85:6 (March 15, 1952), pp. 1038-1039.

Radioactivity of Eu^{152,154}

Robert Katz and Milford R. Lee

Kansas State College, Manhattan, Kansas

Submitted January 28, 1952

Although the spectrographic analysis accompanying several enriched samarium isotopes from Oak Ridge contained <0.6 percent europium, some of the more intense internal conversion lines attributed to the disintegration of europium have been found photographically with neutron-activated samarium isotopes in a 180° focusing β -ray spectrograph.

©1952 The American Physical Society. Used by permission.

Online at URL: <http://link.aps.org/abstract/PR/v85/p1038/s2>

DOI: 10.1103/PhysRev.85.1038.2

Letters to the Editor

Radioactivity of Eu^{152,154}

ROBERT KATZ AND MILFORD R. LEE
Kansas State College, Manhattan, Kansas
(Received January 28, 1952)

ALTHOUGH the spectrographic analysis accompanying several enriched samarium isotopes from Oak Ridge contained ≤ 0.6 percent europium, some of the more intense internal conversion lines attributed to the disintegration of europium¹ have been found photographically with neutron-activated samarium isotopes in a 180° focusing β -ray spectrograph. Conversion lines of some gamma-rays designated by Cork *et al.*¹ as 122.0 (*K, L, M*), 244.3 (*K, L*), and 343.8 (*K, L*) kev were found on spectrograms of neutron-irradiated Sm¹⁵⁰, while conversion lines of these three gamma-rays, together with the *K* conversion line of the 123.2-kev gamma-ray, were found associated with neutron-irradiated Sm¹⁵⁴. Our energies for these gamma-rays were 121.2, 243.8, 343.8, and 122.4 kev, respectively. The spectrograph was calibrated with Cs¹³⁷. Assuming that the europium impurity was enriched in nearly the same way as the samarium, the abundance ratio of Eu¹⁵¹:Eu¹⁵³ would be markedly different in the two samples, being 6:1 in the Sm¹⁵⁰ enrichment and 1:16 in the Sm¹⁵⁴ enrichment. In addition to these long lived conversion electrons, hard gamma-rays (half-life > 1 yr) of energy 990 kev in the Sm¹⁵⁰ sample and 1170 kev in the Sm¹⁵⁴ sample were found by absorption, in lead, in a geometry calibrated with Cs¹³⁷ and Co⁶⁰.

The recently revised gamma-ray assignments in Eu¹⁵² and Eu¹⁵⁴ by Keller and Cork,² insofar as they refer to the transfer of the highest energy (1116 kev) gamma-ray to Eu¹⁵⁴ and of the 344-kev gamma-ray to Eu¹⁵², are in good agreement with our data. These results appear to be in disagreement with Fowler and Schreffer³ who have reported finding coincidences between conversion electrons from the 123- and 344-kev gamma-rays. Our data indicate that the 122-, 244-, and 344-kev gamma-rays are associated with the disintegration of Eu¹⁵² while the 123-kev gamma-ray appears to be associated with the disintegration of Eu¹⁵⁴.

It seems clear that the long lived activity assigned to Sm^{155} by Cork, Fowler, and Schreffler⁴ consisting of gamma-rays of energy 950 and 242 kev (and listed under Sm^{148} in both the latest Seaborg table and National Bureau of Standards circular 499)⁵ was the result of europium contamination. We have found conversion electrons of the 242-kev gamma-ray and/or the "950-kev" gamma-ray (Pb absorber) in nearly the same intensity in pile irradiated Sm^{148} , Sm^{150} , and Sm^{152} . The high energy gamma-ray was considerably weaker in a source of pile-irradiated unenriched samarium, further indicating that this activity was because of an impurity.

¹ Cork, Keller, Rutledge, and Stoddard, Phys. Rev. **77**, 848 (1950).

² H. B. Keller and J. M. Cork, Phys. Rev. **84**, 1079 (1951).

³ C. M. Fowler and R. G. Schreffler, Rev. Sci. Instr. **21**, 740 (1950).

⁴ Cork, Fowler, and Schreffler, Phys. Rev. **74**, 240 (1948).

⁵ K. Way (private communication) stated that this long lived activity was listed as Sm^{148} in National Bureau of Standards Circular 499 (1950), since Sm^{148} was reported to be short lived.
